

§ 558.274

21 CFR Ch. I (4–1–14 Edition)

skin. Avoid contact with skin, eyes, and clothing.

(ii) [Reserved]

[50 FR 33719, Aug. 21, 1985, as amended at 50 FR 42518, Oct. 21, 1985; 51 FR 7397, Mar. 3, 1986; 51 FR 11439, Apr. 3, 1986; 51 FR 14989, Apr. 22, 1986; 51 FR 23737, July 1, 1986; 53 FR 1018, Jan. 15, 1988; 53 FR 11065, Apr. 5, 1988; 54 FR 11519, Mar. 21, 1989; 54 FR 28052, July 5, 1989; 59 FR 51498, Oct. 12, 1994; 61 FR 21076, May 9, 1996; 61 FR 24694, May 16, 1996; 64 FR 42597, Aug. 5, 1999; 65 FR 45712, July 25, 2000; 66 FR 47962, Sept. 17, 2001; 71 FR 27956, May 15, 2006; 79 FR 10982, Feb. 27, 2014]

§ 558.274 Hygromycin B.

(a) *Approvals.* See sponsor numbers in § 510.600(c) of this chapter for Type A medicated articles or Type B medicated feeds as follow:

(1) No. 000986: 2.4 and 8 grams per pound (g/lb).

(2) Nos. 012286 and 051311: 2.4 g/lb.

(3) No. 017790: 0.6 g/lb.

(4) No. 054771: 0.6 and 1.6 g/lb.

(b) *Related tolerances.* See § 556.330 of this chapter.

(c) *Conditions of use.* It is used in feed as follows:

(1) *Chickens—*

Hygromycin B in grams per ton	Combination in grams per ton	Indications for use	Limitations	Sponsor
(i) 8 to 12	Chickens: For control of infestation of large roundworms (<i>Ascaris galli</i>), cecal worms (<i>Heterakis gallinae</i>), and capillary worms (<i>Capillaria obsignata</i>).	Withdraw 3 days before slaughter	000986 012286 017790 054771
(ii) 8 to 12	Tylosin 4 to 50	Chickens: For control of infestations of large roundworms (<i>Ascaris galli</i>), cecal worms (<i>Heterakis gallinae</i>), and capillary worms (<i>Capillaria obsignata</i>); growth promotion and feed efficiency.	Withdraw 3 days before slaughter. Tylosin as tylosin phosphate as provided by No. 000986 in § 510.600 of this chapter.	000986

(2) *Swine—*

Hygromycin B in grams per ton	Combination in grams per ton	Indications for use	Limitations	Sponsor
(i) 12	Swine: For control of infestation of large roundworms (<i>Ascaris suis</i>), nodular worms (<i>Oesophagostomum dentatum</i>), and whipworms (<i>Trichuris suis</i>).	Withdraw 15 days before slaughter.	000986 012286 017790 054771
(ii) 12	Tylosin 10 to 100 ...	Swine: For control of infestations of large roundworms (<i>Ascaris suis</i>), nodular worms (<i>Oesophagostomum dentatum</i>), and whipworms (<i>Trichuris suis</i>); growth promotion and feed efficiency.	Feed continuously as follows: Animal weight (lbs.): Up to 40 . . . 20 to 100 ¹ 41 to 100 . . . 20 to 40 ¹ 101 to market weight . . . 10 to 20 ¹ Withdraw 15 days before slaughter. Tylosin as tylosin phosphate as provided by No. 000986 in § 510.600 of this chapter.	000986

¹ Amount of Tylosin (g/t).

[79 FR 10982, Feb. 27, 2014]

§ 558.295 Iodinated casein.

(a) *Approvals.* See 017762 in § 510.600(c) of this chapter.

(b) *NAS/NRC status.* The use of this drug is NAS/NRC reviewed and found effective. Applications for these uses

need not include efficacy data as required by § 514.111 of this chapter but may require bioequivalency or safety data.

(c) *Conditions of use—*(1) *Ducks—*(i) *Amount per ton.* 100 to 200 grams.